

A filtration defined by arcs on a variety

W.Ebeling and S.M.Gusein-Zade *

Abstract

We define a natural filtration on the ring $\mathcal{O}_{V,0}$ of germs of functions on a germ of a complex analytic variety $(V,0)$ related with the geometry of arcs on the variety and describe some properties of it.

Let $(V,0)$ be a germ of a complex analytic variety and let $\mathcal{O}_{V,0}$ be the ring of germs of functions on it. There is a natural filtration on the ring $\mathcal{O}_{V,0}$ defined by powers of the maximal ideal. We define another natural filtration related with the geometry of arcs on $(V,0)$ and describe some properties of it.

An arc ϕ on $(V,0)$ is a germ of a complex analytic mapping $\phi : (\mathbb{C},0) \rightarrow (V,0)$. For a function $g \in \mathcal{O}_{V,0}$, its order $v_\phi(g)$ on the arc ϕ is defined as the order of the composition $g \circ \phi$, i.e., as the power of the first non-vanishing term in the power series expansion $g \circ \phi(\tau) = a\tau^{v_\phi(g)} + \dots$, $a \neq 0$ (if $g \circ \phi \equiv 0$, then $v_\phi(g)$ is assumed to be equal to $+\infty$). Let $\widehat{v}(g) := \min_{\{\phi\}} v_\phi(g)$. One has $\widehat{v}(g) < \infty$ for $g \neq 0$. One can easily see that $\widehat{v} : \mathcal{O}_{V,0} \setminus \{0\} \rightarrow \mathbb{Z}_{\geq 0}$ is a valuation.

Definition: The *arc filtration* $\mathcal{O}_{V,0} = F_0 \supset F_1 \supset F_2 \supset \dots$ on the ring $\mathcal{O}_{V,0}$ is the filtration by the ideals $F_i = \{g \in \mathcal{O}_{V,0} \mid \widehat{v}(g) \geq i\}$.

On a germ of a smooth variety the arc filtration coincides with the filtration by powers of the maximal ideal.

Suppose that $(V,0)$ has an isolated singularity and let $\pi : (X,D) \rightarrow (V,0)$ be a resolution of it where $D = \bigcup_{i=1}^r E_i$ is a normal crossing divisor and E_i ($i = 1, \dots, r$) are its irreducible components. Let $v_i(g)$ ($g \in \mathcal{O}_{V,0}$) be the

*Partially supported by the DFG-programme "Global methods in complex geometry" (Eb 102/4-2), grants RFBR-01-01-00739, INTAS-00-259. Keywords: ring of functions, arcs on a variety, filtration, Poincaré series. AMS Math. Subject Classification: 14B05, 32S10

order of the lifting $\tilde{g} = g \circ \pi$ of the function g to the space X of the resolution along the component E_i ($i = 1, \dots, r$). This defines a collection of valuations $\{v_1, \dots, v_r\}$ on the ring $\mathcal{O}_{V,0}$ and the corresponding multi-indexed filtration on it: $J(\underline{v}) = \{g \in \mathcal{O}_{V,0} \mid \underline{v}(g) \geq \underline{v}\}$, $\underline{v} = (v_1, \dots, v_r)$, $\underline{v}(g) = (v_1(g), \dots, v_r(g))$ (see, e.g., [CDG3]).

Proposition 1

$$\widehat{v}(g) = \min_{1 \leq i \leq r} v_i(g).$$

Proof. This follows from the fact that the set of arcs on $(V, 0)$ coincides with the set of arcs on (X, D) and the order of the function g along the component E_i of the exceptional divisor D is equal to its order along a generic smooth curve transversal to E_i . $\square$

Let

$$P_{V,0}(t) = \sum_{i=0}^{\infty} \dim(F_i/F_{i+1}) \cdot t^i$$

be the Poincaré series of the arc filtration on the ring $\mathcal{O}_{V,0}$.

In [CDK, CDG3] there was defined a (generalized) Poincaré series $P_{\{v_i\}}(t_1, \dots, t_r)$ corresponding to the collection $\{v_i\}$ of valuations. Strictly speaking there it is defined for the ring of functions on a curve or on a surface, however, there is no difference to the general case.

For a formal power series $P(t_1, \dots, t_r) \in \mathbb{Z}[[\underline{t}]]$ ($\underline{t} = (t_1, \dots, t_r)$), let its *reduction* $\overline{P}(t)$ be the formal power series in one variable t obtained from the series $P(t_1, \dots, t_r)$ by substituting each monomial $\underline{t}^{\underline{v}} = t_1^{v_1} \cdots t_r^{v_r}$ in it by the monomial $t^{\min v_i}$. For example, if $P(t_1, t_2) = 1 + t_1 t_2^2 + t_1^2 t_2$, then $\overline{P}(t) = 1 + 2t$.

Proposition 2 *Let the germ $(V, 0)$ be irreducible. Then the Poincaré series $P_{V,0}(t)$ of the arc filtration coincides with the reduction $\overline{P}_{\{v_i\}}(t)$ of the Poincaré series $P_{\{v_i\}}(t_1, \dots, t_r)$ of the collection $\{v_i\}$ of the divisorial valuations.*

Proof. A convenient way to see this is to express both Poincaré series as certain integrals with respect to the Euler characteristic over the projectivization $\mathbb{P}\mathcal{O}_{V,0}$ of the space $\mathcal{O}_{V,0}$ of germs of functions on $(V, 0)$. The definition of such an integral can be found, e.g., in [CDG2, CDG3]. It is inspired by the notion of motivic integration (see, e.g., [DL]) and in some sense dual to it. Let $\mathbb{Z}[[\underline{t}]]$ be the abelian group (with respect to addition) of formal power series

in $t_1, \dots, t_r$. A monomial $\underline{t}^{v(g)}$ can be considered as a function on the projectivization $\mathbb{P}\mathcal{O}_{V,0}$ of the space $\mathcal{O}_{V,0}$ with values in the group $\mathbb{Z}[[\underline{t}]]$. From [CDG3] it follows that

$$P_{\{v_i\}}(t_1, \dots, t_r) = \int_{\mathbb{P}\mathcal{O}_{V,0}} \underline{t}^{v(g)} d\chi. \quad (1)$$

The proof in [CDG3] is formulated for surface singularities, but it can be easily extended to an arbitrary collection of finitely determined valuations $\{v_i\}$. (A valuation v is *finitely determined* if for any $i \geq 0$ there exists $N \geq 0$ such that $\{g \in \mathcal{O}_{V,0} \mid v(g) \geq i\} \supset \mathfrak{m}^N$.) For the arc valuation $\hat{v}$ it gives

$$P_{V,0}(t) = \int_{\mathbb{P}\mathcal{O}_{V,0}} t^{\hat{v}(g)} d\chi. \quad (2)$$

The formulae (1) and (2) and Proposition 1 imply Proposition 2. $\square$

The statement of Proposition 3 does not hold in general for reducible germs.

Corollary 1 *The reduction $\overline{P_{\{v_i\}}}(t)$ of the Poincaré series $P_{\{v_i\}}(t_1, \dots, t_r)$ of the collection $\{v_i\}$ of the divisorial valuations defined by a resolution does not depend on the resolution. $\square$*

For rational surface singularities, the Poincaré series $P_{\{v_i\}}(t_1, \dots, t_r)$ (for the minimal resolution) were computed in [CDG3]. Using Proposition 2, these results yield the following statement.

Proposition 3 *For the rational double points one has the following list of the Poincaré series $P_{V,0}(t)$ of the arc filtration:*

$$\begin{aligned} A_k &: \frac{1 - t^2}{(1 - t)^3} \\ D_k &: \frac{1 - t^{k-1}}{(1 - t)^2(1 - t^{k-2})} \\ E_6 &: \frac{1 - t^4}{(1 - t)(1 - t^2)^2} \\ E_7 &: \frac{1 - t^6}{(1 - t)(1 - t^2)(1 - t^4)} \\ E_8 &: \frac{1 - t^6}{(1 - t^2)^2(1 - t^3)} \end{aligned}$$

Remarks. 1. The computations, though formal and routine, are somewhat tedious. In each step one has in general expressions which are not products of powers of cyclotomic polynomials. Moreover, for some singularities (e.g., for D_5) they even start from one which is not of this sort (see [CDG3]). However, finally they lead to series which are such products.

2. For an arbitrary singularity $(V, 0)$ the Poincaré series $P_{V,0}(t)$ of the arc filtration must not be a product of powers of cyclotomic polynomials. For example, for the monomial space curve given by $x \mapsto (x^3, x^4, x^5)$ one has $P_{V,0}(t) = (1 - t + t^3)/(1 - t)$. However, for an irreducible *plane* curve singularity the Poincaré series $P_{V,0}(t)$ coincides with the Poincaré series considered, e.g., in [CDG1] and therefore is of this type.

3. Let $(V, 0)$ be an irreducible singularity with a good $\mathbb{C}^*$ -action (i.e., 0 is in the closure of every orbit). Then, if the $\mathbb{C}^*$ -action is free outside of the origin, the arc filtration coincides with the one defined by the natural grading on the ring $\mathcal{O}_{V,0}$ corresponding to the $\mathbb{C}^*$ -action. However, if the action is not free, this is not the case.

4. One can see that for all rational double points the Poincaré series $P_{V,0}(t)$ of the arc filtration (as a rational function of t) has degree -1 and a pole of order 2 at $t = 1$. Moreover, all these series coincide with the Poincaré series with respect to the quasihomogeneous grading of certain rational double points. Namely, this correspondence is the following one:

$$A_k \mapsto A_1, \quad D_k \mapsto A_{k-2}, \quad E_6 \mapsto A_3, \quad E_7 \mapsto A_5, \quad E_8 \mapsto D_4.$$

We don't understand the meaning of this correspondence.

The fact that all singularities A_k have one and the same Poincaré series $P_{V,0}(t)$ is a particular case of the following statement (which can be easily proved).

Proposition 4 *Let f be a germ of a function in n variables $z_1, \dots, z_n$ which belongs to the square $\mathfrak{m}^2$ of the maximal ideal $\mathfrak{m} \subset \mathcal{O}_{\mathbb{C}^n,0}$ and let $(V, 0)$ be the hypersurface singularity in $(\mathbb{C}^{n+2}, 0)$ (with the coordinates $(z_1, \dots, z_n, x, y)$) defined by the equation $f(z_1, \dots, z_n) + x^2 + y^2 = 0$ (the double suspension of the hypersurface singularity $\{f = 0\}$). Then*

$$P_{V,0}(t) = \frac{1 - t^2}{(1 - t)^{n+2}}.$$

The main property which guarantees that $P_{V,0}(t)$ does not depend on f in $\mathfrak{m}^2$ for the hypersurface $V = \{f(z_1, \dots, z_n) + x^2 + y^2 = 0\}$ is that there are many smooth curves on V .

References

- [CDG1] S. M. Gusein-Zade, F. Delgado, A. Campillo: On the monodromy of a plane curve singularity and the Poincaré series of the ring of functions on the curve. *Funct. Anal. Appl.* **33**:1, 56–57 (1999).
- [CDG2] S. M. Gusein-Zade, F. Delgado, A. Campillo: Integration with respect to the Euler characteristic over the projectivization of the space of functions and the Alexander polynomial of a plane curve singularity. *Russian Math. Surveys* **55**, no.6, 1148–1149 (2000).
- [CDG3] A. Campillo, F. Delgado, S. M. Gusein-Zade: Poincaré series of a rational surface singularity. Preprint, University of Valladolid, 2002.
- [CDK] A. Campillo, F. Delgado, K. Kiyek: Gorenstein property and symmetry for one-dimensional local Cohen–Macaulay rings. *Manuscripta math.* **83**, no.3–4, 405–423 (1994).
- [DL] J. Denef, F. Loeser: Germs of arcs on singular algebraic varieties and motivic integration. *Inventiones math.* **135**, no.1, 201–232 (1999).

Universität Hannover, Institut für Mathematik
Postfach 6009, D-30060 Hannover, Germany
E-mail: ebeling@math.uni-hannover.de

Moscow State University, Faculty of Mechanics and Mathematics
Moscow, 119992, Russia
E-mail: sabir@mccme.ru